



April Forecast Update for Atlantic Hurricane Activity in 2018

Issued: 5th April 2018

by Professor Mark Saunders and Dr Adam Lea
 Dept. of Space and Climate Physics, UCL (University College London), UK

Forecast Summary

TSR lowers its forecast and predicts Atlantic hurricane activity in 2018 will be about 15% below the long-term average. However, forecast uncertainties remain large.

The TSR (Tropical Storm Risk) April forecast update anticipates North Atlantic hurricane activity in 2018 will be slightly below-norm. Based on current and projected climate signals, Atlantic basin tropical cyclone activity is forecast to be about 15% below the 1950-2017 long-term norm and 25% below the recent 2008-2017 10-year norm. The forecast spans the period from 1st June to 30th November 2018 and employs data through to the end of March 2018. The main reason why the TSR forecast for North Atlantic hurricane activity in 2018 has fallen by 30% since the extended range outlook issued in December 2017 is the anticipation now for cooler sea surface temperatures during August-September 2018 in the tropical North Atlantic and Caribbean Sea than anticipated previously. However, it should be stressed that the precision of hurricane outlooks issued in early April is low and that forecast uncertainties remain large for the 2018 hurricane season.

Atlantic ACE Index and System Numbers in 2018

		ACE Index	Intense Hurricanes	Hurricanes	Tropical Storms
TSR Forecast	2018	84	2	6	12
68yr Climate Norm	1950-2017	103	3	6	11
10yr Climate Norm	2008-2017	115	3	7	15
Forecast Skill at this Lead	1980-2017	7%	3%	2%	7%

- Key: ACE Index = Accumulated Cyclone Energy Index = Sum of the Squares of 6-hourly Maximum Sustained Wind Speeds (in units of knots) for all Systems while they are at least Tropical Storm Strength. ACE Unit = $\times 10^4$ knots².
- Intense Hurricane = 1 Minute Sustained Wind > 95Kts = Hurricane Category 3 to 5.
 Hurricane = 1 Minute Sustained Wind > 63Kts = Hurricane Category 1 to 5.
 Tropical Storm = 1 Minute Sustained Winds > 33Kts.
 Forecast Skill = Percentage Improvement in Mean Square Error over Running 10-year Prior Climate Norm from Replicated Real Time Forecasts 1980-2017.

There is a 27% probability that the 2018 Atlantic hurricane season ACE index will be above-average (defined as an ACE index value in the upper tercile historically (>119)), a 33% likelihood it will be near-normal (defined as an ACE index value in the middle tercile historically (69 to 119)) and a 40% chance it will be below-normal (defined as an ACE index value in the lower tercile historically (<69)). The 68-year period 1950-2017 is used for climatology.

- Key: Terciles = Data groupings of equal (33.3%) probability corresponding to the upper, middle and lower one-third of values historically (1950-2017).
- Upper Tercile = ACE index value greater than 119.
 Middle Tercile = ACE index value between 69 and 119.
 Lower Tercile = ACE index value less than 69.

ACE Index & Numbers Forming in the MDR, Caribbean Sea and Gulf of Mexico in 2018

		<u>ACE Index</u>	<u>Intense Hurricanes</u>	<u>Hurricanes</u>	<u>Tropical Storms</u>
TSR Forecast	2018	65	2	4	8
68yr Climate Norm	1950-2017	81	2	4	8
10-yr Climate norm	2008-2017	95	3	5	10
Forecast Skill at this Lead	1980-2017	11%	7%	15%	17%

The Atlantic hurricane Main Development Region (MDR) is the region 10°N-20°N, 20°W-60°W between the Cape Verde Islands and the Caribbean Lesser Antilles. A storm is defined as having formed within this region if it reached at least tropical depression status while in the area.

There is a 30% probability that the 2018 Atlantic hurricane season ACE index will be above-average (defined as an ACE index value in the upper tercile historically (>92)), a 35% likelihood it will be near-normal (defined as an ACE index value in the middle tercile historically (43 to 92) and a 35% chance it will be below-normal (defined as an ACE index value in the lower tercile historically (<43)). The 68-year period 1950-2017 is used for climatology.

USA Landfalling ACE Index and Numbers in 2018

		<u>ACE Index</u>	<u>Hurricanes</u>	<u>Tropical Storms</u>
TSR Forecast	2018	1.5	1	2
68yr Climate Norm	1950-2017	2.4	1	3
10yr Climate Norm	2008-2017	2.2	1	3
Forecast Skill at this Lead	1980-2017	0%	1%	4%

Key: ACE Index = Accumulated Cyclone Energy Index = Sum of the Squares of hourly Maximum Sustained Wind Speeds (in units of knots) for all Systems while they are at least Tropical Storm Strength and over the USA Mainland (reduced by a factor of 6). ACE Unit = $\times 10^4$ knots².
 Strike Category = Maximum 1 Minute Sustained Wind of Storm Directly Striking Land.
 USA Mainland = Brownsville (Texas) to Maine

USA landfalling intense hurricanes are not forecast since we have no skill at any lead.

There is a 32% probability that in 2018 the USA landfalling ACE index will be above average (defined as a USA ACE index value in the upper tercile historically (>2.08)), a 25% likelihood it will be near-normal (defined as a USA ACE index value in the middle tercile historically (0.83 to 2.08)) and a 43% chance it will be below-normal (defined as a USA ACE index value in the lower tercile historically (<0.83)). The 68-year period 1950-2017 is used for climatology.

Caribbean Lesser Antilles Landfalling Numbers in 2018

		<u>ACE Index</u>	<u>Intense Hurricanes</u>	<u>Hurricanes</u>	<u>Tropical Storms</u>
TSR Forecast	2018	1.4	0	0	1
68yr Climate Norm	1950-2017	1.5	0	0	1
10yr Climate Norm	2008-2017	1.9	0	1	2
Forecast Skill at this Lead	1980-2017	1%	1%	5%	0%

Key: ACE Index = Accumulated Cyclone Energy Index = Sum of the Squares of hourly Maximum Sustained Wind Speeds (in units of knots) for all Systems while they are at least Tropical Storm Strength and within the region 10°-18°N, 63°-60°W (reduced by a factor of 6). ACE Unit = $\times 10^4$ knots².
 Strike Category = Maximum 1 Minute Sustained Wind of Storm Directly Striking Land.
 Lesser Antilles = Island Arc from Anguilla to Trinidad Inclusive.

Methodology and Key Predictors for 2018

The TSR statistical seasonal hurricane forecast model divides the North Atlantic into three regions and employs separate forecast models for each region before summing the regional hurricane forecasts to obtain an overall forecast. For two of these three regions (tropical North Atlantic, and the Caribbean Sea and Gulf of Mexico) the forecast model pools different environmental fields involving August-September sea surface temperatures (SSTs) and July-September trade wind speed to select the environmental field or combination of fields which gives the highest replicated real-time skill for hurricane activity over the prior 10-year period. The nature of this process means that the details of the seasonal forecast model can vary subtly from year-to-year and also with lead time within the same year. Separate forecast models are employed to predict the July-September trade wind speed and to predict the August-September SSTs. Finally bias corrections are employed for each predictand based on the forecast model performance for that predictand over the prior 10 years.

The main factor underpinning the TSR forecast for 2018 hurricane activity being below the long term norm is the anticipated suppressing effect of the July-September 2018 forecast trade wind at 925mb height over the Caribbean Sea and tropical North Atlantic region (7.5°N – 17.5°N, 100°W – 30°W). The current forecast for this predictor is $0.49 \pm 0.88 \text{ ms}^{-1}$ stronger than normal (1980-2017 climatology). This is notably higher than the December forecast value of $0.37 \pm 0.87 \text{ ms}^{-1}$ weaker than normal. The July-September 2018 trade wind prediction uses the current expectation of neutral ENSO conditions in July-September 2018 as forecast by the consensus of dynamical and statistical model ENSO outlooks (http://iri.columbia.edu/climate/ENSO/currentinfo/SST_table-.html) provided by the International Research Institute for Climate and Society, combined with expected cooler than normal sea surface temperatures in the tropical North Atlantic and Caribbean Sea. Stronger than normal trade winds during July-August-September are associated with less cyclonic vorticity and increased vertical wind shear over the hurricane main development region. These environmental factors reduce hurricane frequency and intensity. However, it should be stressed that uncertainties in the forecast July-September 2018 trade wind speed are large due to the large uncertainties in ENSO and in North Atlantic and Caribbean Sea SSTs at this 4-month lead before the start of the hurricane peak season in August.

The main reason why the TSR forecast for North Atlantic hurricane activity in 2018 has fallen by 30% since the extended range outlook was issued in December 2017 is the anticipated development of cooler than normal sea surface temperatures in the tropical North Atlantic and Caribbean Sea by the summer/autumn of 2018. This decrease in sea surface temperature was not anticipated in December 2017.


The Precision of Seasonal Hurricane Forecasts

The figure on the next page displays the seasonal forecast skill for North Atlantic hurricane activity for the 15-year period between 2003 and 2017. This assessment uses the seasonal forecast values issued publicly in real-time by the three forecast centres TSR, NOAA (National Oceanic and Atmospheric Administration) and CSU (Colorado State University). Skill is assessed as a function of lead time for two measures of hurricane activity: ACE and basin hurricane numbers.

Forecast precision is assessed using the Mean Square Skill Score (MSSS) which is the percentage improvement in mean square error over a climatology forecast. Positive skill indicates that the model performs better than climatology, while a negative skill indicates that it performs worse than climatology. Two different climatologies are used: a fixed 50-year (1951-2000) climatology and a running prior 10-year climate norm.

It should be noted that NOAA does not issue seasonal hurricane outlooks before late May and that CSU stopped providing quantitative extended-range hurricane outlooks from the prior December in 2011. It is clear from the figure that there is little skill in forecasting the upcoming number of hurricanes from the previous December. Skill climbs slowly as the hurricane season approaches with moderate-to-good skill levels being achieved from early August.

TSR was the best performing statistical seasonal forecast model at all lead times for 2003-2017.



Further Information and Next Forecast

Further information about TSR forecasts and verifications may be obtained from the TSR web site <http://www.tropicalstormrisk.com>. The next TSR forecast update for the 2018 North Atlantic hurricane season will be a pre-season forecast issued on the 30th May 2018.

Appendix – Predictions from Previous Months

1. Atlantic ACE Index and System Numbers

Atlantic ACE Index and System Numbers 2018					
		ACE Index	Named Tropical Storms	Hurricanes	Intense Hurricanes
Average Number (1950-2017)		103	11	6	3
Average Number (2008-2017)		115	15	7	3
CSU Forecast	5 Apr 2018	130	14	7	3
TSR Forecasts	5 Apr 2018	84	12	6	2
	7 Dec 2017	117	15	7	3

2. MDR, Caribbean Sea and Gulf of Mexico ACE Index and Numbers

MDR, Caribbean Sea and Gulf of Mexico ACE Index and Numbers 2017					
		ACE Index	Named Tropical Storms	Hurricanes	Intense Hurricanes
Average Number (1950-2017)		81	8	4	2
Average Number (2008-2017)		95	10	5	3
TSR Forecast	5 Apr 2018	65	8	4	2

3. US ACE Index and Landfalling Numbers

US Landfalling Numbers 2018				
		ACE Index	Named Tropical Storms	Hurricanes
Average Number (1950-2017)		2.4	3	1
Average Number (2008-2017)		2.2	3	1
TSR Forecast	5 Apr 2018	1.5	2	1

4. Lesser Antilles ACE Index and Landfalling Numbers

Lesser Antilles Landfalling Numbers 2018					
		ACE Index	Named Tropical Storms	Hurricanes	Intense Hurricanes
Average Number (1950-2017)		1.5	1	0	0
Average Number (2008-2017)		1.9	2	1	0
TSR Forecast	5 Apr 2018	1.4	1	0	0